





























## 5. Outcome Measures

If outcomes are the ultimate indicator of care quality, why did Donabedian propose other approaches to measuring quality of care and why have so many researchers used structure and process measures? One of the reasons is that many outcomes are often not closely linked to the quality of care, in part because outcomes are affected by many social and clinical factors that are not related to the treatment provided. This is particularly true for measures of HRQL (Wilson, I. B. & Cleary, 1995).

**We often do not have the information necessary to identify the most powerful predictors of outcome that are not related to care. Even if we can identify the predictors we are often unsure about how best to specify them.**

One of the major factors affecting post-treatment status is pre-treatment status (Cleary, P. D., Greenfield, et al., 1991). Pre-treatment health may be a predictor of post-treatment health and thus need to be a control variable, or the appropriateness of a procedure may be conditional on health status (e.g. the benefits of a certain type of survey are greater than the potential harm only for healthy patients) and thus need to be considered as a stratifying variable. In many situations, however, "pre-treatment" status is impossible to assess because patients are not identified at the beginning of their illness.

Links between outcome and process are more likely when the patient group is well defined by medical condition and/or demographic characteristics, when there is a well accepted physiologic, biochemical, or psychological mechanism that links medical intervention with outcome, and when the outcomes are targeted for the medical condition.

## 5. Outcome Measures

Another problem is that the outcome of interest may be too rare to give us a reliable picture of quality for the entity of interest. For example, counting breast cancer deaths might seem like a good way to assess the effectiveness of health plans' screening and treatment programs for breast cancer, but only about one of every 1,000 women over the age of fifty die in any given year. Thus, screening (versus no screening) might change a group or clinic's death rate by only three per 10,000 patients a year. It becomes even harder to detect differences when it is not an either/or situation. If rates of screening varied by ten percentage points between clinics or medical groups, the difference in death rates might be only three per 10,000 patients per year (Eddy, 1998). To detect such differences, one would need a very large sample, much larger than might be available for individual providers or even group practices or small health plans. If one wanted to assess the quality of diabetes care using available quality measures, some have estimated that a physician would need to have more than 100 patients with diabetes for the quality measures to have adequate reliability (Hofer, et al., 2000), again a limitation if one wanted to assess the quality of care provided by individual physicians.



## Exercise 3:

Match each of the following examples with the corresponding limitation of outcomes as quality of care measures.

### Limitations:

✚ Outcomes affected by social and clinical factors not related to treatment

✚ Difficult to assess "pre-treatment" status

✚ Outcome of interest too rare

✚ Patient group not well-defined

### Examples:

How long people survive after a heart attack depends in part on life style changes that they make.

How well people can function after hip replacement surgery depends in part on their fitness and mobility prior to their surgery and this is often not measured in a standardized way.

Mortality after gall bladder surgery is very uncommon.

The health status of patients in any given health insurance plan varies greatly.

Patient care patterns are affected by their insurance, which varies greatly among patients.

Stroke patients are typically only identified after they have already had a stroke.



## 5. Outcome Measures

Often we lack information on the natural history of illnesses that are not treated. This would provide an important comparison against which outcomes for treated patients could be compared. In the case of procedures, measuring outcomes can have an unintended and perverse consequence: performing the procedure on healthier patients may give the appearance of better technical quality because the patients' outcomes are better if the case-mix models do not adequately adjust for such differences. If a procedure was not indicated for the patient in the first place, the good outcome may not be attributable to the procedure and the patient may have being exposed to unnecessary risk since the procedure may not have been necessary (Dranove, Kessler, McClellan, & Satterthwaite, 2002).

Other limitations of outcome measures are that in many situations the most relevant outcome takes a substantial amount of time to become manifest. For example, one of the outcomes of a hip replacement might be how long the replacement lasts and it may take many years to have a good sense of that. A limitation of outcome measures for quality improvement purposes is that even when outcomes are less than optimal, they may not provide insights into why there were poor outcomes and what needs to be changed to yield better outcomes.

There are other practical difficulties in assessing outcomes. The data collection systems required to collect such data are not routinely available in clinical settings. When one does develop data collection systems there can be biases in the data. Specifically, for some measures, particularly patient reported ones, patients with worse outcomes may be less likely to respond to surveys. Thus, bias can result if response rates differ across settings.

## 6. Process Measures

Process measures attempt to answer the question “Did this patient receive the right care,” or “what percent of the time did patients of this type receive the right care?” Such measures are typically developed based on the known relationship between a process and outcomes. For example if one was examining the quality of care received by a patient with diabetes, one might assess whether the patient had undergone an annual funduscopy examination by an ophthalmologist or whether the patient's feet were professionally examined annually (Brook, et al., 1996). Such measures are used because research has demonstrated a link between those processes and important outcomes, such as retinopathy or foot amputations. A nurse or medical-record technician trained in quality assessment could compare what was done to what should have been done, and the result would be expressed as the proportion of times that the criteria were met.

Such measures or criteria are typically developed by first identifying the condition of interest, and then synthesizing research evidence to create evidence-based guidelines for clinical care. Once one has identified the part of the medical care process that will be used, one defines patients who are eligible to receive care on the basis of guideline, create criterion to determine which patients received care in accordance with guideline, and divide number who received care in compliance with guideline by number of patients eligible to receive care.

## 6. Process Measures

Researchers increasingly are recognizing that it is not adequate to simply assess individual processes of care, but rather groups or processes, or “bundles” of activities that need to occur to lead to a better outcome. For example, researchers attempting to prevent catheter related bloodstream infections learned from prior research that multiple activities, such as hand washing, full barrier precautions, skin antisepsis with chlorhexidine, avoiding the femoral site during catheter insertion, and removing unnecessary catheters are all necessary to achieve the best outcomes (Lipitz-Snyderman, et al., 2011). Similarly, interventions to prevent ventilator associated pneumonia included a mechanical ventilator “bundle” consisting of use of semirecumbent positioning, daily interruption of sedation infusions, and prophylaxis for peptic ulcer disease and deep venous thrombosis (Berenholtz, et al., 2011).

**An important limitation on the usefulness of process measures is that much care is delivered in the absence of compelling evidence of effectiveness. Although there has been extensive work on the development of evidence based guidelines (Institute of Medicine, 2011), the evidence for many of the specific things that clinicians do is lacking.**

## 6. Process Measures

Even when there are data supporting the appropriateness and effectiveness of a process or procedure, there often is more than one evidence-supported way to treat a condition. Frequently, for example, different approaches to treatment (e.g. radiation, versus surgery for prostate cancer) are developed and thought to be best by physicians in different specialties. This situation has given rise to a field of research referred to as **comparative effectiveness research**.

The goal of **comparative effectiveness research** is to evaluate which of two or more appropriate approaches to treating a condition is better (Institute of Medicine, 2009). That is, although there may be multiple approaches that are considered good quality care, this approach tries to assess which is the best.

It is also important to recognize that for many treatments that are “preference sensitive” whether or not a particular treatment or procedure is appropriate depends on patient preferences (Brook, et al., 1996; Sepucha & Mulley Jr, 2009)

Sometimes, processes of care are too complicated for completely explicit criteria. For example, determining when a problem occurred or when an adverse event was preventable, may require some clinical judgment (Sullivan, et al., 2007). Such measures tend to be less reliable and usually provide less compelling evidence than measures with a strong research base, however. Even with the most careful protocols, there inevitably are many sources of variation in the way implicit criteria are implemented.

## 6. Process Measures

Some of the advantages of process measures are that they are very specific, understandable to patients and providers, and in many cases can be easier than outcomes to measure. They also answer the intuitive question a provider might have: "am I doing the right thing for a patient? A related strength is that they are actionable because they indicate what should be changed. They also can be used to make inferences about individual providers. Disadvantages of process measures include the fact that we do not know how many processes of care are related to outcomes. Another shortcoming is related to the fact that sometimes hundreds, if not thousands of things are done in the course of caring for a patient with a complicated condition and it is difficult to develop and use enough measures to form a comprehensive assessment.

## 7. Structure Measures

Probably the main advantage and attractiveness of structure measures is that they are concrete and usually easy to assess. For example, it is relatively easy to determine whether an intensive care unit has a specialty physician available 24 hours a day or if a health plan provides incentives to physicians who meet high standards of care, the training of physicians (Landon, et al., 2003; Landon, et al., 2002), whether a clinic specializes in particular types of care (Wilson, IB., et al., 2005), or the number of procedures performed per year (Hannan, et al., 1997).

Structural characteristics that did not receive a great deal of attention when Donabedian did his seminal research include organizational culture, including to the priority that a clinic or hospital gives to quality as well as leadership, policies and procedures for maximizing the quality of care (Berwick, 1996; Institute of Medicine, 2001)

The main disadvantage of such measures is that often the association between structure and process and/or structure and outcome are not well-established and developing evidence for such associations is difficult. One of the reasons for this is that the link between structure and process or outcome measures of quality are often very complex (Landon, Wilson, & Cleary, 1998) and consequently weak (Landon, Zaslavsky, Beaulieu, Shaul, & Cleary, 2001). Another weakness is that the most accessible structural variables often lack specificity. For example, one can relatively easily determine if a person is board certified in infectious diseases but it is much harder to develop a measure of the quality of that training or the extent to which the physician uses that knowledge or skills.

## 8. Comprehensiveness of Measures

Available studies indicate that it is difficult to generalize from the quality of care for one set of symptoms or diseases to the quality of care for another set of symptoms or diseases. For example, a study of six medical and surgical interventions at six teaching hospitals showed that the rank ordering of the hospitals' performance in terms of both process and outcome measures differed depending on the

intervention.(Cleary, P. D., Greenfield, et al., 1991). Even for a specific condition, the fact that one aspect of care is performed well may not predict the quality of a different aspect of care for the same condition (Wilson, .IB., et al., 2007). Innovative organizations, such as Intermountain Healthcare and the Boston based Institute for Healthcare Improvement are focusing on improving the systems that should support better care across a wide range of processes, but to date, it does not appear that such comprehensive approaches to quality improvement are widespread enough to allow one to generalize about quality on the basis of a few conditions, diagnoses, or symptoms. Such generalizations are especially problematic when different types of medical functions are evaluated, such as screening, prevention, diagnosis, and treatment. Thus, a sound measure of the quality of care must include separate measures for the care provided by different clinicians or organizations or even different functions or procedures provided by the same clinician.

Unfortunately, developing and using a comprehensive assessment, such as the one used by McGlynn et al. (McGlynn, et al., 2003) is very complicated and expensive. As providers of health care in the United States increasingly adopt electronic medical records and continue standardizing the way information is entered and shared, there undoubtedly will be increasing opportunities to develop more comprehensive process measures of care based on information from those systems.(Bates, Ebell, Gotlieb, Zapp, & Mullins, 2003; Poon, et al., 2010; Simon, S. R., R. Kaushal, P.D. Cleary, C.A. Jenter, L.A. Volk, E.J. Orav, E. Burdick, E.G. Poon, D.W. Bates, 2006; Simon, S. R., et al., 2009)

## 9. Selecting Measures

Given the numerous considerations summarized above, selecting quality of care measures is a daunting task for someone who is not an expert in this type of assessment. However, by specifying different aspects of the measure requirements, one can narrow down the possibilities considerably.

**In our opinion, for the reason summarized above, the assessment of quality should depend much more on process data than on outcome or structure data. Researchers should be aware however, that we recommend the use of measures with a strong evidence base and such evidence is lacking for many process of care that we would like to assess.**

The first task is to specify the study design requirements. A major consideration, for example, is the unit of assessment. Although one usually is assessing the care of individuals, it is important to know if one wants to make decisions about the individual, a clinician, organization, or population (e.g. state). The requirements of such measurement tasks are very different. For example, if one is assessing the care of clinicians, there often is a "small numbers problem," as described above for assessment of the impact of breast cancer screening and diabetes care. That is, a specific clinician may not see enough patients with diabetes during the study period to yield a reliable measure of his or her performance. If one is evaluating the care system, one must determine, for example, whether the focus is hospital care, ambulatory care, or the combined effect of care provides by multiple providers.



## 9. Selecting Measures

There are, of course, exceptions to this, such as the method used to compare differences in outcome after coronary-artery bypass surgery. There has been extensive research on the best way to adjust statistically for case-mix differences when assessing the outcome of such surgery, there is strong evidence of the link between the quality of care and survival, death after coronary-artery bypass is common enough to be used as a measure of differences in the quality of care, and differences in mortality among institutions or among groups of patients receiving coronary-artery bypass surgery can be assessed relatively soon after the surgery. Furthermore, because numerous aspects of the care that are also difficult to measure influence postsurgical mortality (e.g., the physician's skill in the operating room), differences in survival may reflect differences in quality, or case selection, not revealed by a limited number of process assessments and case-mix adjustors. As another example, we might prefer to use adjusted outcomes to evaluate the potential effect of a policy change designed to reduce health care expenditures.

**There recently has been a substantial increase in funding for comparative effectiveness research (Institute of Medicine, 2009). Many comparative effectiveness studies, which will focus on specific procedures in defined populations or systems changes, will appropriately use outcome measures.**

## 9. Selecting Measures

For any of these applications, measures based on explicit criteria derived from strong evidence are more reliable and valid. After the scientific literature has been reviewed, specific criteria of the quality of care are enumerated and categorized by the level of evidence (randomized controlled trials, observational studies, or expert opinion) supporting them. To be useful, these criteria must be as clinically detailed as possible and must cover diverse topics, such as what drugs were used in a patient with asthma, whether smoking cessation was recommended for smokers with stable angina, or whether the appropriate candidates were offered coronary-artery bypass surgery.

## 9. Selecting Measures

After deciding what method of quality assessment should be used, the next step is to determine the appropriate sources of data. Data used in quality assessment are obtained from diverse sources, such as records maintained by insurance companies to reimburse physicians, clinical records maintained by health care professionals, pharmacy and laboratory data, and surveys of patients. Each source of data produces a different view of the quality of care (Wilson, A. & McDonald, 1994).

For example, suppose we asked a patient who had been told she had breast cancer whether the doctor had discussed options for removing the cancer. Because of the emotional impact of the news, the patient might not remember whether the doctor had discussed therapeutic options. The doctor might not have recorded that discussion in the patient's medical record, but an audiotape would have captured the entire conversation.

**The answer depends on the purpose of the assessment. If the purpose is to determine whether the patient comprehended the relevant information, then survey data are most appropriate (Cleary, P., 1999; Cleary, P. D., Edgman-Levitan, et al., 1991; Cleary, P. D. & McNeil, 1988).**

Finally, if one wants to determine whether one doctor recorded enough information to allow another physician caring for the patient to know what had been done, then the medical record might be the best source of data. Thus, the appropriate source of data for quality assessment depends on the purpose for which the information will be used.

## 9. Selecting Measures

Administrative data (e.g. billing data, patient characteristics) are maintained by health care organizations, private insurers, and public insurance programs, such as Medicare and Medicaid. Data often are coded using standardized codes, representing the patient's diagnosis (DRGs; ICD) or the procedures conducted and/or billed for (CPT codes). Limitations of administrative include limited specificity, limited detail, and potential bias. Because reimbursement is often based on diagnosis or procedures performed, there can be incentives to use particular codes, often referred to as "up coding."

There are other types of data such as tumor registry data for all patients with cancer, pharmacy and laboratory data, and hospital discharge data. These types of data can provide information that is not available in administrative data (e.g. very specific details about the type of cancer a patient has) and often they can be linked to administrative data to provide more complete information about the patient and/or process of care.

## 9. Selecting Measures

Medical Records maintained by health care providers usually contain detailed information about a patient's clinical history and current status, and information on test results. Disadvantages are that it usually is very costly to abstract such data from medical records, unless they are in standardized electronic formats. Also documentation can be incomplete and unstandardized. For example, different providers may use different language and characteristics to describe the severity of a patient's condition.

As indicated previously, the expanding use of electronic medical records may transform the way we use medical record information for a variety of purposes, including the assessment of care quality. This is part of a broader social transformation in which enormous quantities of information is being produced in digital form and being made available to different groups and the public at large (King, 2011). Although one still needs to assess carefully the completeness and accuracy of such information, there is the possibility that the standardization of how we collect, store, and transmit such information will fundamentally change the value of such information by making it more accurate, useful, and accessible.

## 9. Selecting Measures

Patient surveys can be used for many purposes in quality studies, but the main uses are to assess outcomes and processes from the patient's perspective, or the extent to which care is "patient centered." The outcomes of treatment are central to patients. As a result, many consider what is generally referred to as Health Related Quality of Life (HRQL), along with survival, key criteria of high quality medical care. In the ambulatory setting, these sorts of data are particularly important because ambulatory medical care is often directed at reducing morbidity more than decreasing mortality.

An advantage of asking patients directly for information about outcomes is that this type of information is less susceptible to "gaming", or biased reporting to improve reimbursement or apparent performance, by providers. Patients certainly can be influenced by their providers and there are biases inherent in the way patients report information but the risk of "gaming" is lower than for information provided by physicians or nurses.



## Exercise 4

The research purpose should drive the decision about what quality of care measures should be used. For each of the following research purposes, decide which measure would be the most appropriate.

	Administrative	Medical records	Patient survey	Provider survey	Patient registry
A researcher wishes to assess whether physicians explain side effects of a prescribed medication to patients in a way that they can understand.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
A researcher wants to study whether African American patients of an appropriate age receive mammograms at the same rate as Caucasian patients.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
A researcher wants to identify patients with cardiac disease that affects the left main artery to study outcomes in this group.	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
A researcher wants to understand if physician awareness and/or agreement with professional society guidelines are related to the extent to which their actions conform with such guidelines	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

**A researcher wants to understand if physician awareness and/or agreement with professional society guidelines are related to the extent to which their actions conform with such guidelines**

**A researcher wants to study the age and sex distribution of lung cancer in Connecticut.**



## **10. Examples of Quality of Care Measures**

The Agency for Health Care Policy and Research has used both literature reviews and expert opinion to establish guidelines for care and quality-of-care criteria. The scientific literature has also been used to develop evidence-based practice guidelines and to evaluate both the appropriateness of use of procedures and the quality of inpatient care received by patients with multiple conditions (Brook, et al., 1996). The Cochrane Centre conducts similar reviews.

The National Quality Forum (NQF) is a nonprofit organization with the mission of improving the quality of health care by: building a consensus on national priorities and goals for performance improvement and working in partnership to achieve them; endorsing national consensus standards for measuring and publicly reporting on performance; and promoting national goals through education and outreach programs.

## **10. Examples of Quality of Care Measures**

The National Committee for Quality Assurance is a private, not-for-profit organization dedicated to improving health care quality. Since its founding in 1990, NCQA has been a central organization in driving health care improvement. One of the ways in which NCQA has pursued its mission is by working with large employers, policymakers, doctors, patients and health plans to decide what aspect of care quality are most important and developing measures of those aspects of care. It has developed quality standards and performance measures for a broad range of health care entities. These measures and standards are the tools that organizations and individuals can use to identify opportunities for improvement. It has developed and implemented a variety of standards for improving health care quality that include structural characteristics of health care organizations, but one of its most widely known measurement sets is the Healthcare Effectiveness Data and Information Set (HEDIS), a set of measures that are used by more than 90 percent of U.S. health plans to measure performance on important dimensions of care and service.

## 12. Resources

- The Patient Reported Outcomes Measurement Information System (PROMIS)
- The Agency for Health Care Policy and Research (AHRQ)
- The Cochrane Centre
- The National Quality Forum (NQF)
- National Committee on Quality Assurance (NCQA)

### 13. References

- Barry, M. J., Fowler, F. J., Jr, Mulley, A. J., Henderson, J. V., & Wennberg, J. E. (1995). Patient reactions to a program designed to facilitate patient participation in treatment decisions for benign prostatic hyperplasia. *Med Care*.
- Bates, D. W., Ebell, M., Gotlieb, E., Zapp, J., & Mullins, H. C. (2003). A proposal for electronic medical records in U.S. primary care. *J Am Med Informatics Assoc*, 10, 1-10.
- Berenholtz, S., Pham, J. C., Thompson, D. A., Needham, D., Lubomski, L., Hyzy, R., et al. (2011). An intervention to reduce ventilator-associated pneumonia in the ICU. *Infect Control Hosp Epidemiol*, In Press.
- Berwick, D. M. (1996). A primer on leading the improvement of systems. *BMJ*, 312, 619-622.
- Bostridge, M. (2008). *Florence Nightingale. The Making of an Icon*. New York, NY: Farrar Straus and Giroux.
- Braddock, C. H., Edwards, K. A., Hasenberg, N. M., & et, a. (1999). Informed decision making in outpatient practice: time to get back to basics. 282, 2313-2320.
- Brook, R. H., McGlynn, E. A., & Cleary, P. D. (1996). Quality of health care. Part 2: measuring quality of care. *N Engl J Med*, 335(13), 966-970.
- Cassedy, J. H. (1984). *American Medicine and Statistical Thinking, 1800-1860*. Cambridge, MA: Harvard University Press.
- Cleary, P. (1999). The increasing importance of patient surveys. *Br Med J*, 319, 720-721.
- Cleary, P. D., Edgman-Levitan, S., Roberts, M., Moloney, T. W., McMullen, W., Walker, J. D., et al. (1991). Patients evaluate their hospital care: a national survey. *Health Aff*, 10(4), 254-267.
- Cleary, P. D., Greenfield, S., Mulley, A. G., Pauker, S. G., Schroeder, S. A., Wexler, L., et al. (1991). Variations in length of stay and outcomes for six medical and surgical conditions in Massachusetts and California. *JAMA*, 266(1), 73-79.
- Cleary, P. D., & McNeil, B. J. (1988). Patient satisfaction as an indicator of quality care. *Inquiry*, 25(1), 25-36.

Codman, E. A. (1914). The product of a hospital. *Surg, Gynecol, Obstet*, 18, 491-496.

Deyo, R. A., Diehr, P., & Patrick, D. L. (1991). Reproducibility and responsiveness of health status measures statistics and strategies for evaluation. *Controlled Clinical Trials*, 12(4, Supplement 1), S142-S158.

Donabedian, A. (1965). Evaluating the quality of medical care. *Milbank Memorial Fund Quart*, 44(3), 166-206.

Donabedian, A. (1980). *Explorations in quality assessment and monitoring. Vol. 1. The definition of quality and approaches to its assessment*. Ann Arbor, Mich.: Health Administration Press.

Donabedian, A. (1982). *Explorations in quality assessment and monitoring. Vol. 2. The criteria and standards of quality*. Ann Arbor, Mich.: Health Administration Press.

Donabedian, A. (1985). *Explorations in quality assessment and monitoring. Vol. 3. The methods and findings of quality assessment and monitoring: an illustrated analysis*. Ann Arbor, Mich.: Health Administration Press.

Donabedian, A. (1988). The quality of care. How can it be assessed? *JAMA*, 260(12), 1743-1748.

Dranove, D., Kessler, D., McClellan, M., & Satterthwaite, M. (2002). *Is More Information Better? The Effects of 'Report Cards' on Health Care Providers*. Boston, MA: NBER.

Eddy, D. M. (1998). Performance measurement: problems and solutions. *Health Aff*, 17(4), 7-25.

Hannan, E. L., Racz, M., Ryan, T. J., McCallister, B. D., Johnson, L. W., Arani, D. T., et al. (1997). Coronary angioplasty volume-outcome relationships for hospitals and cardiologists. *JAMA*, 277(11), 892-898.

Hofer, T. P., Hayward, R. A., Greenfield, S., Wagner, E. H., Kaplan, S. H., & Manning, W. G. (2000). The unreliability of individual physician "report cards" for assessing the costs and quality of care of a chronic disease. *JAMA*, 281(22), 2098-2105.

Institute of Medicine. (2000). *To Err is Human*. Washington D.C.: National Academy Press.

Institute of Medicine. (2001). *Crossing the Quality Chasm: A New Health System for the 21st Century*. Washington D.C.: National Academy Press.

Institute of Medicine. (2009). *Initial National Priorities for Comparative Effectiveness Research*. Washington, DC: The National Academies Press.

Institute of Medicine. (2011). *Clinical Practice Guidelines We Can Trust*. Washington, DC: The National Academies Press.

King, G. (2011). Ensuring the data-rich future of the social sciences. *Science*, 331(11 Feb.).

Landon, B. E., Wilson, I. B., & Cleary, P. D. (1998). A conceptual model of the effects of health care organizations on the quality of medical care. *JAMA*, 279(17), 1377-1382.

Landon, B. E., Wilson, I. B., Cohn, S. E., Fichtenbaum, C. J., Wong, M. D., Wenger, N. S., et al. (2003). Physician specialization and antiretroviral therapy for HIV. *J Gen Intern Med*, 18(4), 233-241.

Landon, B. E., Wilson, I. B., Wenger, N. S., Cohn, S. E., Fichtenbaum, C. J., Bozzette, S., et al. (2002). Specialty training and specialization among physicians who treat HIV/ AIDS in the United States. *J Gen Intern Med*, 17, 12-22.

Landon, B. E., Zaslavsky, A. M., Beaulieu, N. D., Shaul, J. A., & Cleary, P. D. (2001). Health plan characteristics and consumers' assessments of quality. *Health Affairs*, 20(2), 274-286.

Leape, L. (1994). Error in medicine. *JAMA*, 272, 1851-1857.

Lipitz-Snyderman, A., Steinwachs, D., Needham, D. M., Colantuoni, E., Morlock, L., & Pronovost,

P. J. (2011). Impact of a statewide intensive care unit quality improvement initiative on hospital mortality and length of stay: retrospective comparative analysis. 342(d219).

Lohr, K., & Committee to Design a Strategy for Quality Review and Assurance in Medicare (Eds.). (1990). *Medicare: a strategy for quality assurance, Vol. 1*. Washington, DC: IOM, National Academy Press.

McDowell, I., & Newell, C. (1996). *Measuring Health: A Guide to Rating Scales and Questionnaires*. New York: Oxford University Press.

McGlynn, E. A., Asch, S. M., Adams, J., Keesey, J., Hicks, J., DeCristofaro, A., et al. (2003). The quality of health care delivered to adults in the United States. *N Engl J Med*, 348(26), 2635-2645.

Neuhauser, D. (1990). Ernest Amory Codman, M.D., and end results of medical care. *Int J Tech Assess Health Care*, 6, 307-325.

Poon, E. G., Wright, A., Simon, S. R., Jenter, C. A., Kaushal, R., Volk, L. A., et al. (2010). Relationship between use of electronic health record features and health care quality: Results of a Statewide Survey. *Med Care*, 48(3), 203-209.

President's Commission for the Study of Ethical Problems in Medicine and Biomedical and Behavioral Research. (1982). *Making Health Care Decisions*. Wash, DC: Gov't Printing Office.

Schuster, M. A., McGlynn, E. A., & Brook, R. H. (1998). How good is the quality of health care in the United States? *Milbank Q*, 76(4), 517-563.

Sepucha, K., & Mulley Jr, A. G. (2009). A perspective on the patient's role in treatment decisions. *Med Care Res Rev*, 66, 53S-74S.

Simon, S. R., R. Kaushal, P.D. Cleary, C.A. Jenter, L.A. Volk, E.J. Orav, E. Burdick, E.G. Poon, D.W. Bates. (2006). How and Why Are Physicians Using Electronic Health Records? A Statewide Survey. *Arch Int Med*.

Simon, S. R., Soran, C. S., Kaushal, R., Jenter, C. A., Volk, L. A., Burdick, E., et al. (2009). Physicians' usage of key functions in electronic health records from 2005 to 2007: A statewide survey. *J Am Med Informatics Assoc*, 16(4), 465-470.

Stewart, A. L., Hays, R. D., & Ware, J. E. J. (1988). The MOS short-form General Health Survey (SF-36) 1. Conceptual framework and item selection. *Med Care*, 26, 724-735.

Sullivan, A. F., Camargo, C. A., Cleary, P. D., Gordon, J. A., Guadagnoli, E., Kaushal, R., et al. (2007). The National Emergency Department Safety Study: Study rationale and design. *Acad Emerg Med*, 14(2), 1182-1189.

Terwee, C. B., Dekker, F. W., Wiersinga, W. M., Prummel, M. F., & Bossuyt, P. M. M. (2003). On Assessing Responsiveness of Health-Related Quality of Life Instruments: Guidelines for Instrument Evaluation. *Quality of Life Research*, 12(4), 349-362.

Tsevat, J., Weeks, J. C., Guadagnoli, E., Tosteson, A. N., Mangione, C. M., Pliskin, J. S., et al. (1994). Using health-related quality-of-life information: clinical encounters, clinical trials, and health policy. *J Gen Intern Med, 9*(10), 576-582.

Ware, J. J., Kosinski, M., & Keller, S. (1996). A 12-item short-form health survey. Construction of scales and preliminary tests of reliability and validity. *Med Care, 34*(3), 220-223.

Wilson, A., & McDonald, P. (1994). Comparison of patient questionnaire, medical record, and audio tape in assessment of health promotion in general practice consultations. *BMJ, 309*, 1483-1485.

Wilson, I., Landon, B., Ding, L., Zaslavsky, A. M., Shapiro, M. F., Bozzette, S. A., et al. (2005). A national study of the relationship of care site HIV specialization to early adoption of highly active antiretroviral therapy. *Med Care, 43*(1), 12-20.

Wilson, I., Landon, B. E., Marsden, P. V., Hirschhorn, L. R., McInnes, K., Ding, L., et al. (2007). Correlations among quality measures in HIV Care in the United States: a cross-sectional study of care sites in 30 states. *Br Med J, 335*(7629), 1085-1091.

Wilson, I. B., & Cleary, P. D. (1995). Linking clinical variables with health-related quality of life. A conceptual model of patient outcomes. *JAMA, 273*(1), 59-65.



## 14. Author Biographies

**Paul D. Cleary, Ph.D.** is the Anna M.R. Lauder Professor of Public Health and Dean of the Yale School of Public Health. Dr. Cleary received his Ph.D. in sociology from the University of Wisconsin. He has been a member of the Institute of Medicine since 1994 and a member of the Connecticut Academy for Science and Engineering since 2007. In 1996, he was selected as a distinguished fellow of the Association for Health Services Research, and in 2002, received the Distinguished Investigator Award from the Academy for Health Services Research and Health Policy.

Dr. Cleary's research includes a study of how organizational characteristics affect the costs and quality of care for persons with AIDS; a national evaluation of a continuous quality improvement initiative in clinics providing care to HIV infected individuals; and a study of the long-term impact of patient-centered hospital care. He is Principal Investigator of one of the Consumer Assessment of Healthcare Providers and Systems (CAHPS) projects funded by the Agency for Healthcare Research and Quality (AHRQ) to develop surveys for consumers regarding their health plans and services. He also is Principal Investigator and Director of the Yale Center for Interdisciplinary Research on AIDS (CIRA).

**Margaret E. O'Kane** has served as President of the National Committee for Quality Assurance (NCQA), an independent, nonprofit organization that improves health care quality through measurement, transparency and accountability, since 1990.

Under O'Kane's leadership, NCQA developed the first national system for measuring health plans' quality performance, the first and most rigorous system for accrediting health plans, and the most widely adopted program for transforming medical practices into patient-centered medical homes.

O'Kane has served as co-chair of the National Priorities Partnership and is a board member of the Foundation for Informed Decision Making and the American Board of Medical Specialties. She was elected a member of the Institute of Medicine in 1999 and received the 2009 Picker Institute Individual Award for Excellence in the Advancement of Patient-Centered Care. In 2011, *Modern Healthcare* named O'Kane one of the Top 25 Women in Healthcare.